Malette et al.		
[54]	METHOD OF ACHIEVING HEMOSTASIS, INHIBITING FIBROPLASIA, AND PROMOTING TISSUE REGENERATION IN A TISSUE WOUND	
[76]	Inventors:	William G. Malette, 667 Parkwood La., Omaha, Nebr. 68132; Herbert J. Quigley, 9511 Mockingbird Dr., Omaha, Nebr. 68127
[*]	Notice:	The portion of the term of this patent subsequent to Jul. 19, 2000 has been disclaimed.
[21]	Appl. No.:	440,039
[22]	Filed:	Nov. 8, 1982
Related U.S. Application Data		
[63]	Continuation-in-part of Ser. No. 251,321; Apr. 6, 1981; Pat. No. 4,394,373.	
[51] [52]	Int. Cl. ³	
[58]	Field of Sea	rch
[56]		References Cited
	U.S. F	PATENT DOCUMENTS

United States Patent [19]

[11] Patent Number:

4,532,134

[45] Date of Patent:

* Jul. 30, 1985

3,911,116 10/1975 Balassa 424/95

OTHER PUBLICATIONS

Balassa et al., Chem. Abst., vol. 91, (1979), p. 13962u. Prudden et al., Chem. Abst., vol. 73, (1970), p. 43828a. Prudden et al., The Amer. J. of Surgery, vol. 119, May 1970, pp. 560–564.

Primary Examiner—Sam Rosen
Attorney, Agent, or Firm—Zarley, McKee, Thomte,
Voorhees & Sease

[57] ABSTRACT

The method of achieving hemostasis, inhibiting fibroplasia, and promoting tissue regeneration is described wherein a chitosan solution, or water-soluble chitosan in various solid forms is placed in contact with the tissue wound. The chitosan forms a coagulum which prevents bleeding and negates the formation of a blood clot thereby preventing the formation of fibrin strands. The prevention of the formation of fibrin strands prevents the proliferation of fibroblasts and the synthesis of collagen thereby allowing the promotion of normal tissue regeneration. The process described hereinabove results in wound healing with minimal scar formation. The use of chitosan in grafts is contemplated as well as the use of polyglucosamine in wound healing.

16 Claims, 1 Drawing Figure